

Salmonella and Campylobacter **in broiler transport cages**

Mark Berrang
USDA-Agricultural Research Service
Russell Research Center



FSIS Public Meeting 2/23/06



Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent





Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent





Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Pathogens and transport coops

- *Salmonella* has been detected in coops
- U. S. study: *Salmonella* found on 8/160 (5%) coops prior to use and on 16/160 (10%) coops after use
Bailey et al., JFP 2002 64:1690-1697
- Belgian study: 56/128 (43.8%) crates positive prior to use
Heyndrickx et al, E&I 2002 129:253-265
 - Showed prevalence of *Salmonella* + broilers increased due to transport...authors suggest that crates are important source
- Similar suggestions have been made relative to *Campylobacter*

Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Source of *Campylobacter* and *Salmonella*

- *Campylobacter* and *Salmonella* are present in the gut and feces of broilers from positive flocks
- Feed withdrawal prior to catching can result in change in bacterial profile of alimentary tract
- Stress associated with transport leads to higher rates of excretion which continues during holding
- The result is that *Campylobacter* and *Salmonella* can contaminate surfaces in transport coops

Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Coop flooring and carcass microbiology

Buhr et al 2000 PS 79:436-441

- Study compared broilers transported on solid floors to those transported on elevated wire floors (less contact with feces)
- Feathered carcasses transported on solid flooring had noticeably more fecal contamination...higher numbers of *E. coli*
- However, after defeathering all differences in broiler carcass microbiology disappeared

Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Cross contamination

Berrang et al, 2003 JAPR 12:190-195

- A study was conducted to examine the possibility that contaminated feces left in a dump coop can cause transfer of *Campylobacter* to broilers that were previously free of Campy
- Fecal contamination was isolated as the only source of *Campylobacter* to the test broilers

Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Results

- After just 2 hr exposure to contaminated feces, more than half the defeathered carcasses from test broilers (*Campylobacter* free flock) had become positive for *Campylobacter*
- Numbers on test carcasses were lower than on positive control carcasses (*Campylobacter* positive flock), but still substantial

Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Conclusions

- *Campylobacter* can be spread to previously negative broilers by contact with contaminated feces remaining in a dump coop
- Such contamination can remain on the carcass through scalding and picking.
- Similar findings have been reported with *Salmonella* Rigby et al, 1982 Can J Comp Med 46:272-278

Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent





Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Is washing and sanitizing the answer?

- About 28 % of US broiler plants have a coop wash or sanitize procedure in use
Northcutt and Jones 2004 JAPR 13:48-54
- A higher percentage of processors in the EU wash/sanitize crates
- Nevertheless, European studies report detection of *Salmonella* and *Campylobacter* on crates even after washing and sanitizing

Slader et al 2002 AEM 68:713-719; Corry et al 2002 JAM 92:424-432

Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Washing and sanitizing studies

- Reports exist in the literature that show some experimental washing and sanitizing procedures can work
- Research needs to be evaluated carefully
 - How were the chemicals inactivated before culture?
(are the target bacteria being inadvertently killed after sample collection?)
 - How realistic or commercially viable is the procedure?
 - How expensive or unwieldy is the equipment?
(immersion tanks and high temperature application may be expensive)

Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Coop washing test at commercial processing plant

Northcutt and Berrang

- Measured numbers of total aerobic bacteria and *E. coli* as well as presence of *Salmonella* and *Campylobacter*
- Washing coops by spraying with water reduced numbers of aerobic bacteria and *E. coli*.
- Application of sanitizer lowered numbers more
- Prevalence of *Campylobacter* and *Salmonella* was lessened
- However, *Campylobacter* was still detected in 2/27 (7%) coop floor samples after washing and sanitizing

Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent




Novel approaches: Allowing coops to dry between uses

Berrang et al., 2004 PS 83:1213-1217


- Objective: determine the effect of storing dump coops between uses on the numbers of viable *Campylobacter* in feces that was deposited on the floor by *Campylobacter* positive broilers
- Hypothesis: allowing feces to dry out and remain exposed to atmospheric oxygen would lower the numbers of *Campylobacter* detected in feces

Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent






Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent




Mean Log₁₀ *Campylobacter* detected per opening (all feces)



Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Percent moisture of feces collected from coop at time intervals



Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Conclusions

- Allowing transport coops to dry for 24–48h between uses results in lower numbers of *Campylobacter* remaining in the coops
- Such drying should not be counted on to eliminate *Campylobacter* entirely
- The expense required to maintain enough coops for extended dry times would be hard to justify

Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent




Washing and sanitizing coop flooring

Berrang and Northcutt, 2005 JAPR 14:315-321

- Study designed to test the efficacy of spray washing and immersion in chemical sanitizers to eliminate *Campylobacter* on coop floor material.
- A more controlled study than the field study which tested washing and sanitizing under commercial conditions

Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent





Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent






Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent




Water spray wash followed by 15 s immersion



Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent




Water spray wash followed by 60 s immersion



Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Water spray wash followed by 5 min. immersion



Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Summary and conclusions

- Spraying the floor surface with tap water lowered the numbers of *Campylobacter* recovered.
- Adding an immersion in 200 ppm QAC or chlorine did not help
- The re-moistening effect may have actually revived Campy making recovery more likely

Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Water spray followed by drying as sanitizing step instead of chemical


Berrang and Northcutt, 2005 PS 84:1797-1801

- Measure effectiveness of water spray followed by an extended dry time to lower numbers of *Campylobacter* on soiled coop flooring
- Examine the effect of re-wetting on numbers of *Campylobacter* in gut contents previously allowed to dry on coop floor surface

Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Effect of water spray with and without subsequent drying time (log cfu/square)



* When rewet, *E. coli* numbers rebounded to 2.0

Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Water spray and drying

- Low pressure tap water spray lowered numbers of *Campylobacter* and *E. coli* on soiled coop flooring
- Simply allowing contaminated gut contents to dry out on the flooring had a greater effect on numbers of bacteria recovered
- Spray washing followed by drying was very effective in lowering bacterial numbers
- Re-moistening dried on gut contents caused an rebound in numbers of *E. coli* (*Salmonella*??) but not *Campylobacter*

Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Floor surface drying as a sanitation treatment?

- Allowing floor surface to thoroughly dry between uses could be part of an effective strategy to lower *Campylobacter* contamination in transport coops
- Would require a change in thinking relative to coop design and/or management
- Unsure how *Salmonella* would be affected, more research is required to determine role of VNC *Campylobacter* in treated coop

Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent



Overall

- Coop washing and sanitizing is an expensive proposition (water costs, personnel and time)
- Questionable efficacy
- New and innovative procedures may be more effective than traditional methods
- Ongoing research towards the goal of making coop washing practical, effective and affordable

Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent





Bacterial Epidemiology and Antimicrobial Resistance Research Unit
Not for public distribution without consent

